



Nova™ Fungicide

GROUP	3	FUNGICIDE
-------	---	-----------

In Water-Soluble Pouches

COMMERCIAL (AGRICULTURAL)

SOLUBLE GRANULES

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Myclobutanil 40%

REGISTRATION NO. 22399 PEST CONTROL PRODUCTS ACT

CAUTION  **POISON**

WARNING – EYE IRRITANT

DO NOT SELL THE WATER SOLUBLE PACKETS INDIVIDUALLY

NET CONTENTS: 0.56 kg (4 x 140 g Water-Soluble Pouches)

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS
KEEP OUT OF REACH OF CHILDREN
DO NOT APPLY BY AIR

Hazardous to humans and domestic animals. Causes eye irritation. A potential skin sensitizer. May cause irritation to the nose, throat and skin. Harmful if swallowed, inhaled, or absorbed through the skin. **DO NOT** get in eyes, on skin or breathe spray mist.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Keep the following personal protective equipment immediately available for use in case of emergency (i.e., a broken package, spill, or equipment breakdown): chemical-resistant coveralls, chemical-resistant gloves, chemical-resistant foot wear, chemical-resistant head gear and a respirator.

Use only properly calibrated groundboom, chemigation or hand held equipment as specified by the label.

Not for use by homeowners or other uncertified users.

DO NOT use in residential areas. Residential areas are defined as sites where bystanders including children may be potentially exposed during or after spraying. This includes around homes, school, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed.

Personal Protective Equipment

Wear goggles, mid-forearm to elbow-length chemical-resistant gloves, chemical-resistant footwear, a wide brimmed hat, chemical-resistant coveralls over long pants and a long-sleeved shirt and an appropriate respirator when mixing, loading, and applying this product. Pants or coveralls should be worn outside footwear to prevent pooling within boots.

Remove protective equipment immediately after handling this product. Wash outside of gloves and footwear before removing. As soon as possible, wash thoroughly and change into clean clothing. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Contaminated clothing must be laundered separately in hot water before reusing. Wash hands and face thoroughly after handling and before eating, drinking, chewing gum, smoking, or using toilet.

DO NOT enter treated areas for a minimum of 12 hours for all crops, unless a longer REI is specified. Wear gloves, long sleeved shirts, long pants, a hat and work boots when entering treated areas, including greenhouses, for harvesting, pruning, thinning, suckering or for any other agricultural practice in the treated area. **DO NOT** apply this product in such a manner as to directly, or through drift, expose workers or other persons. Unprotected persons must be vacated from the area being treated. Only protected handlers may be in the area during application.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to birds and small wild mammals. **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**. The use of this product may result in contamination of groundwater particularly in areas where soils are permeable (for example, sandy soil) and/or the depth to the water table is shallow. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

DO NOT allow effluent or runoff from greenhouses or mushroom houses containing this product to enter lakes, streams, ponds or other waters.

STORAGE

Store in a cool dry place but not below 0 deg C. Do not remove package from container except for immediate use. Store this product away from food or feed.

DISPOSAL

1. When all the water-soluble bags are used, make the empty outer package unsuitable for further use.
2. Dispose of the outer package in accordance with provincial/ territorial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Nova Fungicide is systemic fungicide with both after-infection (eradicator) and protectant properties. This fungicide is most effective when used in a routinely scheduled preventative spray program.

GENERAL USE PRECAUTIONS

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

DO NOT apply by air.

Nova Fungicide is a soluble granule sealed within a water soluble bag. **DO NOT** open or puncture water soluble bag for any reason. **DO NOT** use opened or punctured water soluble bag for any reason. If broken water soluble bags are found when container is opened, avoid contact with, and inhalation of the product. Wear chemical resistant coveralls, chemical resistant gloves and a respirator to dispose of broken water soluble bags according to **DISPOSAL** section.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

DIRECTIONS FOR USE

READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC FUNGICIDE CONTROL OR CROP DAMAGE.

Observe a 120-day plant-back interval between the last application and planting leafy vegetable or grain crops, and a 210-day plant-back interval between the last application and planting root vegetable or any other unlabelled crop at the treatment site.

Mixing

The enclosed pouches of Nova Fungicide are water-soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately five minutes from the time they are added to the water. Copper products tank mixed with Nova Fungicide reduce the effectiveness of the fungicide.

Compatibility

Please note - Do not add pouches directly to diesel oils or summer type oil. The pouches are water soluble not oil soluble. Spray tank must be completely free of oil residue from previous sprays. Do not add pouches to spray solutions containing boron or that release free chlorine as the plastic in the pouches becomes insoluble.

Application

A uniform spray coverage is essential for good disease control. Nova Fungicide is a systemic fungicide and will not be affected if rainfall occurs at least one hour after application. This fungicide is locally systemic.

Apply at the recommended rate using a maximum spray volume of 1000 L per hectare, unless otherwise stated.

Restricted Entry Intervals

Crop	Activity	REI* (days)
apples, cherries (sweet & sour), peaches, nectarines	thinning	12
	hand harvest	5
	hand pruning, scouting, pinching, tying, training, hand weeding, propping, animal control, mechanical harvest (cherries only)	0.5
asparagus	All	2
grapes	cane turning and girdling	14
	hand harvesting & pruning, training, thinning, tying, leaf pulling	7
	hand line irrigation, scouting, hand weeding	0.5
strawberries	hand harvest, pinching, pruning, training	2
	irrigation, mulching, scouting, hand weeding	0.5
Saskatoon berries	hand harvest, hand pruning, hand thinning	3
	scouting, hedging, irrigating, hand weeding	0.5
carnations	all	17
Outdoor ornamental trees & shrubs: pear (flowering), crab apple,(flowering), privet, dogwood, euonymus, hawthorn, juniper (flowering & non-flowering), honeysuckle, lilac, crab-apple (flowering); nursery crops: ash, amelanchier	all	0.5
Outdoor ornamental roses	all	11
Outdoor ornamental flowers, shrubs: iris, roses, hollyhock, phlox	all	12
Kentucky bluegrass grown for seed	harvesting/transplanting treated turf, mowing, watering, irrigation, aerating, fertilizing, hand pruning, mechanical weeding, scouting, seeding	0.5
Greenhouse (potted) roses, gerbera, aster, chrysanthemums, geraniums, poinsettias	All	0.5
Greenhouse (cut) roses, gerbera, aster, chrysanthemums, geraniums,	All	1
Greenhouse peppers, tomatoes, cucumbers, eggplant	All	0.5

Crop	Activity	REI* (days)
Nursery (potted) roses, gerbera, aster, chrysanthemums,	All	0.5
Nursery (cut) roses, gerbera, aster, chrysanthemums,	All	12

*Day at which the dermal exposure results in an MOE ≥ 100 or the minimum label REI of 1 or 0.5 days.

DIRECTIONS FOR USE

Apples		
Best control of labeled diseases is achieved when Nova Fungicide is applied on a 7 to 10 day application schedule.		
Disease	Rate of Nova (g/ha)	Use Recommendations
SCAB Prebloom	340	Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Nova Fungicide alone or in a tank mix with 3 kg per hectare Dithane™ Rainshield Fungicide
Bloom and Postbloom	340	Use Nova Fungicide in a tank mix with 3 kg per hectare Dithane DG or Dithane M-45 Rainshield Fungicide for improved fruit scab and summer disease control.
Post- infection (Prebloom only)	340	Nova Fungicide provides 96-hour after-infection control or curative activity. Apply Nova Fungicide as soon as possible after the infection period and follow with a second Nova Fungicide application 7 days later. Continue with a regular Nova Fungicide spray schedule or a regular protectant program. Nova Fungicide will suppress sporulation in established lesions when used in two or more applications.
POWDERY MILDEW	340	Begin application at tight cluster and continue through the second (apples only) cover spray. If powdery mildew was prevalent the previous year, apply Nova Fungicide beginning at green tip. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure.
CEDAR APPLE RUST & QUINCE RUST	340	Begin applications at pink stage and continue through the second cover spray.
RESTRICTIONS Use a maximum of 6 applications per year with a minimum of 7 days between applications. Do not apply Nova Fungicide within 14 days of harvest. Check the labels of Dithane Rainshield Fungicide for preharvest intervals.		

Grapes		
Disease	Rate of Nova (g/ha)	Use Recommendations
POWDERY MILDEW	200	Begin application at prebloom and continue application on a 21 day schedule.
BLACK ROT	200	Begin application at prebloom and continue application on a 14 day schedule during infection periods
RESTRICTIONS Use a maximum of 5 applications per year with a minimum of 14 days between applications. Do not apply Nova Fungicide within 14 days of harvest.		

Stonefruit		
Disease	Rate of Nova (g/ha)	Use Recommendations
CHERRIES - SWEET		
BROWN ROT	340	Begin application when the fruit blossoms are opening. Continue fungicide application on a 7 to 10 day schedule until harvest.
CHERRIES - SOUR		
BROWN ROT POWDERY MILDEW LEAF SPOT	340	Begin application at bloom. Repeat applications at 10 day intervals. Where leaf spot and powdery mildew are serious problems apply further post harvest sprays.

Peaches and Nectarines		
Disease	Rate of Nova (g/ha)	Use Recommendations
BROWN ROT	340	Begin application when the first blossoms are opening and repeat at 7 to 14 day intervals. If wet weather occurs, repeat applications every 4 - 5 days until shuck fall. Protect ripening fruit with fungicide applications starting 2 - 3 weeks before harvesting.
POWDERY MILDEW	340	Begin application after bloom schedule at shuck fall. Repeat at 2 - 4 week intervals until terminal growth ceases.
RESTRICTIONS Use a maximum of 6 applications per year with a minimum of 7 days between applications. Do not apply within 1 day of harvest. Rotate recommended fungicides to help prevent resistance development.		

Saskatoon Berry		
Disease	Rate of Nova (g/ha)	Use Recommendations
POWDERY MILDEW	11.3 g product per 100 L water	Apply at flowering, small green berries, and at green berry stage; spray until dripping.
RESTRICTIONS Do not apply more than 340 g/ha. Use a maximum of 3 applications per year with a minimum of 14 days between applications. Do not apply Nova Fungicide within 14 days of harvest.		

Greenhouse Tomatoes		
Disease	Rate of Nova (g/ha)	Use Recommendations
POWDERY MILDEW	340	Use a maximum of one application per crop cycle. Apply as soon as possible after initial infection; apply 1000 L water/ha; do not apply 3 days before harvest.

Greenhouse Cucumbers		
Disease	Rate of Nova (g/ha)	Use Recommendations
POWDERY MILDEW and GUMMY STEM BLIGHT	340	Use a maximum of one application per crop cycle. Apply when disease first appears. Do not apply 2 days before harvest.

Kentucky Bluegrass Grown for Seed		
Disease	Rate of Nova (g/ha)	Use Recommendations
POWDERY MILDEW (Erysiphe graminis)	250	Apply as a foliar spray. Apply a maximum of two applications, with a minimum of 14 days between applications, one at pre-heading (15-20 cm tall and not yet headed) and again at 50-100% heading.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: The **DIRECTIONS FOR USE** for the uses described in this section of the label were developed by persons other than Corteva Agriscience Canada Company under the User Requested Minor Use Label Expansion program. For these uses, Corteva Agriscience Canada Company has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Refer to the main Nova Fungicide label for additional details and instructions, including preharvest intervals, before using.

READ THE ROTATIONAL CROPPING RESTRICTIONS ON THE FULL LABEL BEFORE USING THIS PRODUCT.

CROP	DISEASE	RATE	RECOMMENDATIONS	RESTRICTIONS
Geraniums	Rust (<i>Puccinia pelargonii-zonatis</i>)	340 g/1000 L water	Apply when disease appears and then every 10-14 days	Use a maximum of 6 applications per growing season, with 10 days between applications.
Poinsettias	Powdery Mildew	280 g/1000 L water	Apply when disease first appears and then every 10-14 days	Use a maximum of 6 applications per growing season, with 10 days between applications.
Asparagus	Rust	340 g/ha	Post harvest treatment of ferns; ground; 370 L water/ha	Use a maximum of 5 applications at 7 day intervals
Cut and Potted Roses, Gerbera, Aster and Chrysanthemums (Nursery and Greenhouse)	Powdery Mildew (<i>Sphaerotheca pannosa</i>); (<i>Erysiphe cichoracearum</i>) and Rust	340 g/1000 L water	Apply as soon as possible after initial infection. Apply as a high volume wet spray to runoff.	Use a maximum of 6 applications per year (Nursery and Greenhouse potted), Greenhouse cut; use a maximum of 4 applications per crop cycle. 10 days between applications.
Nursery Grown Ornamentals:				
Amelanchier Ash Chrysanthemum Iris Hollyhock Phlox	Rust	250-340 g/1000 L water	Apply as soon as possible after initial infection. Apply as a high volume wet spray to runoff.	Use a maximum of 6 applications per year with 10 days between applications.
Carnation	Rust	540 g/1000 L water		
Hollyhock Phlox Rose	Powdery mildew	340 g/1000 L water		
Crabapple Flowering	Rust, Powdery mildew and Scab	340 g/1000 L water		
Greenhouse Peppers	Powdery Mildew	340 g/ha	Apply as soon as possible after initial infection. Apply 1500-3000 L water/ha.	Use a maximum of one application per crop cycle. Do not apply Nova Fungicide within 3 days of harvest.
Crabapple, flowering Hawthorn, flowering Juniper Pear, flowering	Rust	340 g/1000 L water	Apply as soon as possible after initial infection. Apply as a high volume wet spray to runoff.	Use a maximum of 6 applications per year, with 14 days between applications. 10 days between applications for crabapple.

Azalea/Rhododendron Dogwood Euonymus Honeysuckle Lilac Privet	Powdery mildew	340 g/1000 L water		
Dogwood	Anthracnose and Septoria leaf spot	340 g/1000 L water		
Rose	Black spot	340 g/1000 L water	Apply as soon as possible after initial infection. Apply as a high volume wet spray to runoff.	Use a maximum of 4 applications per year with 10 days between applications.
Sweet cherry	Powdery mildew	340 g/ha	Apply 340 g Nova Fungicide/ha at bloom and repeat at 10-14 day intervals as required until terminal growth ceases. Apply 1000 L water/ha.	Use a maximum of 6 applications per year with 7 days between applications. Application at husk fall is important for fruit protection. Pre-harvest interval is 1 day. Application is by ground only.
Strawberry	Powdery mildew	340 g/ha	Begin applications when disease first appears or when conditions favour disease development. Repeat application at 14-to-21 day intervals.	Use a maximum of 6 applications per growing season with 14 days between applications. Do not apply Nova Fungicide within 3 days of harvest.
Pear	Scab (<i>Venturia pirina</i>)	340 g/ha	Apply starting at greentip or when conditions are favorable for scab infestation. Repeat application at 7-to-10 day intervals. Apply as a foliar spray in a spray volume of 500 to 1000 L/water.	Use a maximum of 6 applications per growing season with 14 days between applications. Do not apply within 14 days of harvest.
	Powdery mildew (<i>Podosphaera leucotricha</i>)	340 g/ha	Begin applications at pink stage and repeat at petal fall stage. Repeat application at 7 to 10 day intervals. Coverage to 4 weeks after bloom may be required on highly susceptible varieties. Apply as a foliar spray in a spray volume of 500 to 1000 L/water.	Use a maximum of 6 applications per growing season with 14 days between applications. Do not apply within 14 days of harvest.

Cucurbit Vegetables (Crop Group 9) (Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd, balsam apple and pear, Chinese cucumber, cantaloupe, muskmelon, pumpkin, summer and winter squash, watermelon)	Powdery mildew <i>(Podosphaera xanthii)</i>	175 g/ha	Apply at the first sign of disease development and again 10-14 days later. Apply in a minimum of 250 L water/ha.	Use a maximum of 2 applications per year. Do not apply Nova Fungicide within 3 days of harvest.
Bushberry 13-07B Subgroup (including highbush blueberry, lowbush blueberry, currant, gooseberry)	Powdery Mildew <i>(Microsphaera vaccinii-blueberry)</i> <i>(Spaerotheca sp.-gooseberry and currant)</i>	340 g/ha	Apply at the first sign of disease development and repeat in 7-14 days. Pre-bloom, bloom, and immediately after bloom are the most critical times for the control of powdery mildew and white pine blister rust. Apply with a minimum of 250 L water/ha.	Do not apply more than 340 g/ha. Use a maximum of 3 applications per growing season. Preharvest interval when harvesting by hand is 6 days except for lowbush blueberries, which is 4 days. Preharvest interval when harvesting mechanically is 1 day.
Currant, gooseberry	White Pine Blister Rust <i>(Cronartium ribicola)</i>			
Caneberry 13-07A Subgroup (including raspberry and blackberry)	Powdery Mildew <i>(Spaerotheca sp)</i>	340 g/ha	Apply at the first sign of disease development and repeat in 7-14 days. Pre-bloom, bloom, and immediately after bloom are the most critical times for the control of powdery mildew. Apply with a minimum of 250 L water/ha.	Do not apply more than 340 g/ha. Use a maximum of 3 applications per growing season. Preharvest interval when harvesting by hand is 6 days. Preharvest interval when harvesting mechanically is 1 day.

	Yellow Rust (<i>Phragmidium spp.</i>)	175 g/ha	Begin application as early as bud break when new growth is 3-4 inches or at first sign of yellow rust pustules. Apply with a minimum of 250 L water/ha. Apply at 10-14 day intervals. Use the shorter spray interval under heavy disease pressure. Consult with a crop management specialist for any additional local use recommendations.	
Grape	Suppression of Anthracnose (<i>Elsinoe ampelina</i>)	340 g/ha	Apply as a foliar spray in sufficient water to insure adequate coverage. Begin application when new shoots are 3 to 8 cm in length and continue on an application schedule. Repeat application at 14 day intervals.	Use a maximum of 5 applications per growing season with a minimum of 14 days between applications. Do not apply within 14 days of harvest.
Dry Beans (<i>Phaseolus vulgaris</i>)	Control of Rust (<i>Uromyces appendiculatus</i> . syn. <i>U. phaseoli</i>)	340 g/ha	Apply as soon as weather conditions are favourable for rust development or when first rust pustules are present and continue if conditions remain favourable. Apply as a foliar spray in 300 litres of water to insure thorough coverage of all plant foliage.	Use a maximum of 3 applications per growing season with a 7-10 day re-treatment interval. Do not apply within 30 days of harvest. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 2 days for scouting and 12 hours for all other activities.

Greenhouse Eggplant	Powdery mildew (<i>Leveillula taurica</i> ; <i>Oidium neolycopersici</i>)	340 g/ha	Begin application as soon as possible after initial infection. Maximum 1 application per crop cycle. Apply as a foliar spray in a maximum spray volume of 1000 L water/ha.	Pre harvest interval: 3 days
Globe Artichoke	Powdery mildew (<i>Erysiphe cichoracearum</i> and <i>Leveillula taurica</i>)	280 g/ha	Begin applications when disease first appears or when conditions favour disease development. Repeat application at 14 day intervals. For best results use a minimum of 1000 L of spray volume.	Use a maximum of 6 applications per growing season with 14 days between applications. Pre harvest interval: 3 days

SPRAY BUFFER ZONES

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				Terrestrial Habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Kentucky bluegrass for seed production		0	0	1	0	0
	Strawberry Bushberry 13-07B Subgroup: highbush blueberry, lowbush blueberry, currant, gooseberry Caneberry 13-07A subgroup: raspberry and blackberry dry bean, asparagus juniper shrub Nursery cut and potted ornamentals: rose, gerbera, aster, chrysanthemum, carnation Small nursery grown ornamentals (low plants and small shrubs): rose, iris, hollyhock, phlox, geranium, poinsettias, azalea/rhododendron, dogwood, euonymus, honeysuckle, lilac, privet Globe artichoke		1	0	1	1	1
Airblast	Bushberry 13-07B Subgroup: highbush blueberry, currant, gooseberry Caneberry 13-07A subgroup: raspberry, blackberry Saskatoon berry (amelanchier)	Early growth stage	2	0	2	1	1
		Late growth stage	1	0	1	1	1
	Juniper tree	Early growth stage	2	0	3	1	1
		Late growth stage	1	0	2	1	1

	Grapes	Early growth stage	3	0	4	1	2
		Late growth stage	2	0	2	1	1
	Apples, peaches, nectarines, cherries (sweet and sour), pears	Early growth stage	4	0	5	2	2
		Late growth stage	2	0	3	1	1
	Nursery grown ornamental (tall shrubs): rose, azalea/rhododendron, euonymus, honeysuckle, lilac, privet						
	Nursery grown ornamental (trees): Saskatoon berry, ash, dogwood, flowering crabapple, flowering hawthorn, flowering juniper, flowering pear						

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Nova Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Nova Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of Nova Fungicide or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Corteva Agriscience Canada Company if reduced sensitivity of the pathogen to Nova Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

™Trademark of Corteva Agriscience and its affiliated companies.

181224

Label Code: CN-22399-016-E

Replaces: CN-22399-015-E

Specimen Label Notes

Update to tank mix statement and change address to Quarry Park