

Cirpreme™ XC

with Arylex™ active

Your worst annual AND perennial broadleaf weeds don't stand a chance.

HERBICIDE

WHY USE CIRPREME™ XC HERBICIDE?

- Exceptional control of BOTH annual AND perennial broadleaf weeds in wheat and barley.
- Just GO benefit of Arylex™ active. Consistent, reliable performance across a wide range of staging and environmental conditions.
- Two different modes of action provide overlapping control of certain key broadleaf weeds.

CIRPREME™ XC HERBICIDE ALONE (REQUIRES A NON-IONIC SURFACTANT)

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Canada fleabane
- Chickweed
- Cleavers
- Cow cockle
- Flixweed
- Lamb's-quarters
- Narrow-leaved

- hawk's beard
- Redroot pigweed
- Round-leaved mallow
- Scentless chamomile
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's-bill
- Velvetleaf
- Volunteer canola

- Volunteer flax
- Volunteer sunflower
- Wild buckwheat
- Wild mustard

GRASS WEEDS CONTROLLED

- Barnyard grass

PERENNIAL WEEDS CONTROLLED

- Canada thistle
- Dandelion

WEEDS SUPPRESSED

- Hemp-nettle
- Kochia²
- Night-flowering catchfly
- Perennial sow thistle
- White cockle

CIRPREME XC + 235 ML/AC MCPA ESTER 600 (5 OZ/AC)

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sow thistle*
- Annual sunflower*
- Burdock
- Canada fleabane*
- Chickweed*
- Cleavers*
- Cocklebur
- Cow cockle*
- Field horsetail³ (top growth)
- Flixweed*

- Hemp-nettle
- Henbit
- Lamb's-quarters¹
- Narrow-leaved hawk's beard*
- Plantain³ (top growth)
- Prickly lettuce
- Redroot pigweed*
- Round-leaved mallow*
- Russian pigweed*
- Scentless chamomile
- Shepherd's purse*
- Smartweed*
- Stinkweed*

- Stork's-bill*
- Velvetleaf
- Vetch
- Volunteer alfalfa
- Volunteer canola (all HT biotypes)
- Volunteer flax
- Volunteer sunflower*
- Wild buckwheat*
- Wild mustard*
- Wild radish

- Volunteer flax
- Volunteer sunflower*
- Wild buckwheat*
- Wild mustard*
- Wild radish

GRASS WEEDS CONTROLLED

- Barnyard grass

PERENNIAL WEEDS CONTROLLED

- Canada thistle
- Dandelion
- Perennial sow thistle

WEEDS SUPPRESSED

- Kochia²
- Night-flowering catchfly
- White cockle

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING	• Cirpreme A: 1 x 0.8 kg jug • Lontrel™ XC: 1 x 4.1 L jug
	RATES	• Cirpreme A: 10 g/ac • Lontrel XC: 51 ml/ac
	ACRES TREATED	80 ac/case
	TIMING	Crop stage: 3-leaf to just prior to flag leaf emergence
	RAINFAST	4 hours
	TANK MIX HERBICIDES	• Simplicity™ • Simplicity GoDRI™ • Axial® • Everest®



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

The year following application, fields can be seeded to:

- Barley
- Canola
- Flax
- Forage grasses
- Oats
- Mustard
- Peas⁴
- Soybeans⁴
- Wheat

GRAZING AND HARVEST

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

For a complete weed list and specific weed staging, please refer to the Cirpreme XC label

*Weed controlled through multi-effective mode of action

¹ Including Group 2 resistant biotypes

² Light to moderate infestations (up to 150 plants/m²; up to 15 cm in height), including Group 2 resistant biotypes

³ Cirpreme XC + 6 oz/ac of MCPA Ester 600

⁴ For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).