

Exhilarate™

Arylex™ active

HERBICIDE

With Arylex™ active, Just GO benefits, controls all your standard broadleaf weeds in wheat and barley.



WHY USE EXHILARATE™ HERBICIDE?

- Delivers performance and value for farmers looking to control their most common standard weeds in wheat and barley.
- Contains Arylex™ active which allows farmers to **Just GO** on small or large weeds, early or late crop staging and even in cool or dry conditions.
- Two different modes of action provide overlapping control on certain key broadleaf weeds.

EXHILARATE A + 189 ml/ac PLUS M (4 oz/ac OF MCPA ESTER 600)

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sow thistle
- Canada fleabane*¹
- Chickweed*²
- Cleavers*²
- Common ragweed¹
- Cow cockle*
- Dandelion
- Flixweed
- Henbit
- Hemp-nettle²
- Lamb's-quarters²
- Narrow-leaved hawk's beard
- Redroot pigweed
- Round-leaved mallow*
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's-bill
- Velvetleaf
- Volunteer alfalfa
- Volunteer canola (all herbicide tolerant biotypes)
- Volunteer flax
- Wild buckwheat
- Wild mustard

GRASS WEEDS CONTROLLED

- Barnyard grass

WEEDS SUPPRESSED

- Canada thistle
- Kochia²
- Night-flowering catchfly
- Perennial sow thistle
- Scentless chamomile
- White cockle

*Weed controlled through multi-effective mode of action

² Including Group 2 resistant biotypes

¹ Including Group 2 and Group 9 resistant biotypes

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING	• Exhilarate A: 1 x 0.8 kg jug • Plus M (MCPA Ester 600): 2 x 7.56 L
	RATES	• Exhilarate A: 10 g/ac • Plus M: 189 ml/ac
	ACRES TREATED	80 ac/case
	WATER VOLUME	• Ground 20-40 L/ac (5-10 US gal/ac) • Aerial not registered
	TIMING	Crop stage: 3-leaf to just prior to flag leaf emergence
	RAINFAST	1 hour
	TANK MIX HERBICIDES	• Simplicity™ • Simplicity GoDRI™ • Axial® • Everest®



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Alfalfa
- Barley
- Canola
- Corn
- Dry bean
- Faba beans
- Flax
- Field peas
- Potatoes (except seed potatoes)
- Juncea canola
- Oriental, brown and yellow mustard
- Soybeans
- Oats
- Spring wheat
- Sunflower

GRAZING AND PRE-HARVEST INTERVAL

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.