

Rexade™

Arylex™ active

HERBICIDE

With Arylex™ active Just GO benefits, controls all your standard broadleaf weeds in wheat.



WHY USE REXADE™ HERBICIDE IN WHEAT?

- No compromise. Unmatched cross-spectrum grass and broadleaf weed control in one product.
- New class of Group 4 broadleaf herbicide combined with a premium Group 2 grass herbicide.
- Convenience. Complete, all-in-one box solution.
- GoDRI™ Rapid Dispersion Technology for easy storage, transport and mixing.
- Multi-effective mode of action control of key weeds for effective resistance management.

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sunflower
- Bluebur
- Burdock
- Canada fleabane
- Chickweed²
- Cleavers³
- Cocklebur
- Common ragweed⁴
- Corn spurry
- Cow cockle²
- Flixweed²
- Hemp-nettle^{2,3}
- Henbit⁴
- Kochia^{3,1}
- Lamb's-quarters^{2,3}

- Mustard² (except dog and green tansy)
- Plantain
- Prickly lettuce
- Redroot pigweed^{2,3}
- Round-leaved mallow²
- Russian thistle¹
- Shepherd's purse²
- Smartweed
- Stinkweed²
- Stork's-bill
- Sweet clover
- Velvetleaf
- Volunteer alfalfa
- Volunteer canola
- Volunteer flax
- White cockle
- Wild buckwheat²
- Wild radish¹

GRASS WEEDS CONTROLLED

- Wild oats
- Barnyard grass
- Japanese brome
- Yellow foxtail

WEEDS SUPPRESSED

- Annual sow thistle
- Canada thistle
- Dandelion
- Downy brome
- Green foxtail
- Night-flowering catchfly

1,2 to 4-leaf staging
2 Weed controlled through multi-effective mode of action

3 Including Group 2 resistant biotypes
4 Up to bud stage and 15 cm in height
5 Including Group 2 and Group 9 resistant biotypes

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING <ul style="list-style-type: none"> • Rexade A: 1 x 1.62 kg jug • Rexade B: 1 x 8.58 L jug
	RATES <ul style="list-style-type: none"> • Rexade A: 40.5 g/ac • Rexade B: 215 ml/ac
	ACRES TREATED <p>40 ac/case</p>
	TIMING <p>Crop stage: 2-leaf to flag leaf</p>
	RAINFAST <p>2 hours</p>

MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 to 3/4 of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of Rexade A
4. Add the required amount of Rexade B
5. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler
No tank mix purchase necessary, everything included in the box

CROP ROTATION

All major crops the following spring, except lentils or chickpeas.

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Livestock may graze treated areas 3 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.